



2101 South IH-35, Suite 105
Austin, Texas 78741
(512) 916-4100
(512) 916-4102

November 13, 2018

Reference: Cold Stress Guide

- 1) Anyone working in a cold environment may be at risk of cold stress. Some workers may be required to work outdoors in cold environments and for extended periods, for example, security officers, sanitation workers, police officers and emergency response and recovery personnel, like firefighters, and emergency medical technicians. Cold stress can be encountered in these types of work environment. The following frequently asked questions will help workers understand what cold stress is, how it may affect their health and safety, and how it can be prevented.

How cold is too cold?

- 2) What constitutes extreme cold and its effects can vary across different areas of the country. In regions that are not used to winter weather, near freezing temperatures are considered "extreme cold." A cold environment forces the body to work harder to maintain its temperature. Whenever temperatures drop below normal and wind speed increases, heat can leave your body more rapidly.
- 3) Wind chill is the temperature your body feels when air temperature and wind speed are combined. For example, when the air temperature is 40°F, and the wind speed is 35 mph, the effect on the exposed skin is as if the air temperature was 28°F.
- 4) Cold stress occurs by driving down the skin temperature and eventually the internal body temperature (core temperature). This may lead to serious health problems, and may cause tissue damage, and possibly death.
- 5) What are the risk factors that contribute to cold stress?
 - a) Some of the risk factors that contribute to cold stress are:
 - b) Wetness/dampness, dressing improperly, and exhaustion
 - c) Predisposing health conditions such as hypertension, hypothyroidism, and diabetes
 - d) Poor physical conditioning
- 6) How does the body react to cold conditions?
 - a) In a cold environment, most of the body's energy is used to keep the internal core temperature warm. Over time, the body will begin to shift blood flow from the



2101 South IH-35, Suite 105
Austin, Texas 78741
(512) 916-4100
(512) 916-4102

extremities (hands, feet, arms, and legs) and outer skin to the core (chest and abdomen). This shift allows the exposed skin and the extremities to cool rapidly and increases the risk of frostbite and hypothermia. Combine this scenario with exposure to a wet environment, and trench foot may also be a problem.

7) What are the most common cold induced illnesses/injuries?

- a) Hypothermia
- b) Frostbite
- c) Trench Foot

8) What is hypothermia?

- a) *Hypothermia occurs* when body heat is lost faster than it can be replaced and the normal body temperature (98.6°F) drops to less than 95°F. Hypothermia is most likely at very cold temperatures, but it can occur even at cool temperatures (above 40°F), if a person becomes chilled from rain, sweat, or submersion in cold water.

9) What are the symptoms of hypothermia?

- a) Mild symptoms:
 - i) An exposed worker is alert.
 - ii) He or she may begin to shiver and stomp the feet in order to generate heat.
- b) Moderate to Severe symptoms:
 - i) As the body temperature continues to fall, symptoms will worsen and shivering will stop.
 - ii) The worker may lose coordination and fumble with items in the hand, become confused and disoriented
 - iii) He or she may be unable to walk or stand, pupils become dilated, pulse and breathing become slowed, and loss of consciousness can occur. A person could die if help is not received immediately.

10) What can be done for a person suffering from hypothermia?



2101 South IH-35, Suite 105
Austin, Texas 78741
(512) 916-4100
(512) 916-4102

- a) Call 911 immediately in an emergency; otherwise seek medical assistance as soon as possible.
- b) Move the person to a warm, dry area.
- c) Remove wet clothes and replace with dry clothes, cover the body (including the head and neck) with layers of blankets; and with a vapor barrier (e.g. tarp, garbage bag). Do **not** cover the face.
- d) If medical help is more than 30 minutes away:
 - i) Give warm sweetened drinks if alert (no alcohol), to help increase the body temperature. Never try to give a drink to an unconscious person.
 - ii) Place warm bottles or hot packs in armpits, sides of chest, and groin. Call 911 for additional rewarming instructions.
- e) If a person is not breathing or has no pulse:
 - i) Call 911 for emergency medical assistance immediately.
 - ii) Treat the worker as per instructions for hypothermia, but be very careful and do not try to give an unconscious person fluids.
 - iii) Check him/her for signs of breathing and for a pulse. Check for 60 seconds.
 - iv) If after 60 seconds the affected worker is not breathing and does not have a pulse, trained workers may start rescue breaths for 3 minutes.
 - v) Recheck for breathing and pulse, check for 60 seconds.
 - vi) If the worker is still not breathing and has no pulse, continue rescue breathing.
 - vii) Only start chest compressions per the direction of the 911 operator or emergency medical services*
 - viii) Reassess patient's physical status periodically.
- f) *Chest compression are recommended only if the patient will not receive medical care within 3 hours.



2101 South IH-35, Suite 105
Austin, Texas 78741
(512) 916-4100
(512) 916-4102

11) What is frostbite?

- a) *Frostbite* is an injury to the body that is caused by freezing of the skin and underlying tissues. The lower the temperature, the more quickly frostbite will occur. Frostbite typically affects the extremities, particularly the feet and hands. Amputation may be required in severe cases.

12) What are the symptoms of frostbite?

- a) Reddened skin develops gray/white patches.
- b) Numbness in the affected part.
- c) Feels firm or hard.
- d) Blisters may occur in the affected part, in severe cases.

13) What can be done for a person suffering from frostbite?

- a) Follow the recommendations described above for hypothermia.
- b) Do not rub the affected area to warm it because this action can cause more damage.
- c) Do not apply snow/water. Do not break blisters.
- d) Loosely cover and protect the area from contact.
- e) Do not try to rewarm the frostbitten area before getting medical help; for example, do not place in warm water. If a frostbitten area is rewarmed and gets frozen again, more tissue damage will occur. It is safer for the frostbitten area to be rewarmed by medical professionals.
- f) Give warm sweetened drinks, if the person is alert. Avoid drinks with alcohol.

14) What is immersion/trench foot?

- a) *Trench Foot* or immersion foot is caused by prolonged exposure to wet and cold temperatures. It can occur at temperatures as high as 60°F if the feet are constantly wet. Non-freezing injury occurs because wet feet lose heat 25-times faster than dry feet. To prevent heat loss, the body constricts the blood vessels to shut down circulation in the feet. The skin tissue begins to die because of a lack of oxygen and nutrients and due to the buildup of toxic products.



2101 South IH-35, Suite 105
Austin, Texas 78741
(512) 916-4100
(512) 916-4102

15) What are the symptoms of trench foot?

- a) Redness of the skin, swelling, numbness, blisters

16) What can be done for a person suffering from immersion foot?

- a) Call 911 immediately in an emergency; otherwise seek medical assistance as soon as possible.
- b) Remove the shoes, or boots, and wet socks.
- c) Dry the feet.